

**Adam Seidl, MD**  
Assistant Professor  
Shoulder/Elbow Surgery  
Department of Orthopedics  
University of Colorado



**Orthopedics**  
UNIVERSITY OF COLORADO

**Highlands Ranch Hospital**  
(720) 516-4090  
**Inverness**  
(303) 694-3333  
**Anschutz Medical Campus**  
(720) 848-1900

## Rehabilitation Protocol Revision Reverse Shoulder Arthroplasty

### **Phase I: Protect the Replacement** (*0 to 6 weeks*)

- Sling/immobilizer should be worn AT ALL TIMES for 2 weeks then when sleeping and out in public
- Initiate exercise program 3 times per day immediately:  
    Immediate elbow, forearm and hand AROM
- Pendulums may begin at 2 weeks after instruction by therapist
- At 2 weeks begin PASSIVE supine ROM to 130 FE and 30 ER

### **Phase II: PROM/AAROM** (*6 to 8 weeks*)

- Discontinue sling at all times
- Lifting restriction of 2-3 pounds
- Advance AAROM and PROM as tolerated  
    Maintain ER limit of 30 until 8 weeks.  
    Advance forward elevation as tolerated
- Scapular stabilizer strengthening.

### **Phase III: AROM/Strengthening** (*>8 weeks*)

- Advance AROM as tolerated
- Strengthen rotator cuff and shoulder musculature (Isometrics, Theraband, dumbbell, etc). AVOID RESISTED IR OR EXTENSION UNTIL 10 WEEKS.
- Lifting restriction of 10 pounds until 3 months
- Incorporate low level functional activities at 3 months (swimming, water aerobics, light tennis, jogging)
- Start higher level activities at 4 months (tennis, light weight training, and golf).
- Initiate functional progression to sports specific activities at 4 months.